

An Inequality for the Multivariate Normal Distribution*

LOUIS H. Y. CHEN

*National University of Singapore, Singapore, and
Massachusetts Institute of Technology*

Communicated by T. L. Lai

Herman Chernoff used Hermite polynomials to prove an inequality for the normal distribution. This inequality is useful in solving a variation of the classical isoperimetric problem which, in turn, is relevant to data compression in the theory of element identification. As the inequality is of interest in itself, we prove a multivariate generalization of it using a different argument.

1. INTRODUCTION

Let X be a random variable having the standard normal distribution and let g be an absolutely continuous real-valued function defined on the real line such that $g(X)$ has finite variance. In [2], Chernoff used Hermite polynomials to prove that

$$\text{Var}[g(X)] \leq E[g'(X)]^2 \quad (1.1)$$

and that equality holds if and only if $g(x) = ax + b$ for some constants a and b . As is mentioned in [2], this interesting inequality is useful in solving a variation of the classical isoperimetric problem which, in turn, is relevant to data compression in the theory of element identification (see also Chernoff [1]).

In this paper, we prove a multivariate generalization of inequality (1.1) using a different argument. In our results the finite variance assumption is dropped.

Received October 10, 1980; revised November 1, 1981.

AMS 1980 subject classifications: Primary 60E05, 60E15.

Key words and phrases: Inequality, multivariate normal distribution.

* Research supported in part by National Science Foundation Grant MCS78-01108.

2. THE MAIN THEOREM.

We shall adopt the convention that $\int_0^x = -\int_x^0$ if x is negative. The Lebesgue measure in \mathbb{R}^i will be denoted by μ_i for $i \geq 1$. We first prove a lemma which is itself a slight extension of Chernoff's inequality in the univariate case.

LEMMA 2.1 *Let X be a random variable having the standard normal distribution. Let ψ and ψ' be functions defined on the real line such that ψ is equal a.e. w.r.t. μ_1 to an indefinite integral of ψ' . Then*

$$\text{Var}[\psi(X)] \leq E \left[\int_0^x \psi'(t) dt \right]^2 \leq E[\psi'(X)]^2. \quad (2.0)$$

Furthermore, $\text{Var}[\psi(X)] = E[\psi'(X)]^2 < \infty$ if and only if $\psi(x) = ax + b$ a.e. w.r.t. μ_1 for some constants a and b .

Proof. Let

$$f(x, t) = I(x > t \geq 0) + I(x < t < 0),$$

where $I(A)$ denotes the indicator function of the set A . Then

$$\left(\int_0^x \psi'(t) dt \right)^2 = \left(\int_{-\infty}^{\infty} f(x, t) \psi'(t) dt \right)^2$$

and by the Cauchy-Schwarz inequality

$$\begin{aligned} \left(\int_0^x \psi'(t) dt \right)^2 &\leq \left(\int_{-\infty}^{\infty} f(x, t) dt \right) \left(\int_{-\infty}^{\infty} f(x, t) [\psi'(t)]^2 dt \right) \\ &= |x| \int_{-\infty}^{\infty} f(x, t) [\psi'(t)]^2 dt \end{aligned}$$

where it is noted that $f^2 = f$. Therefore,

$$\begin{aligned} E \left(\int_0^x \psi'(t) dt \right)^2 &\leq E |X| \int_{-\infty}^{\infty} f(X, t) [\psi'(t)]^2 dt \\ &= \int_{-\infty}^{\infty} [E |X| f(X, t)] [\psi'(t)]^2 dt \\ &= \int_{-\infty}^{\infty} (2\pi)^{-1/2} e^{-t^2/2} [\psi'(t)]^2 dt \\ &= E[\psi'(X)]^2. \end{aligned}$$

This proves the second inequality in (2.0). The first inequality in (2.0) follows from the assumption that $\psi(x)$ differs from $\int_0^x \psi'(t) dt$ by a constant for almost all x w.r.t. μ_1 .

If $\psi(x) = ax + b$ a.e. w.r.t. μ_1 for some constants a and b , then trivially $\text{Var}[\psi(X)] = E[\psi'(X)]^2 < \infty$. Conversely, suppose the latter holds. Then using the fact that the standard normal distribution is equivalent to μ_1 , we have

$$\left(\int_{-\infty}^{\infty} f(x, t) \psi'(t) dt \right)^2 = \left(\int_{-\infty}^{\infty} f(x, t) dt \right) \left(\int_{-\infty}^{\infty} f(x, t) [\psi'(t)]^2 dt \right)$$

for almost all x w.r.t. μ_1 . This implies that for almost all x w.r.t. μ_1 $f(x, t) \psi'(t) = a(x) f(x, t)$ or $f(x, t) = 0$ for almost all t w.r.t. μ_1 , where $a(x)$ depends only on x . It is not difficult to see that the second alternative that $f(x, t) = 0$ for almost all t w.r.t. μ_1 is not possible for x belonging to a set of positive measure. So for almost all x w.r.t. μ_1 we have $\psi'(t) = a(x)$ for almost all t w.r.t. μ_1 such that $0 < t \leq x$ or $t < x < 0$. This implies that for almost all t w.r.t. μ_1 ,

$$\begin{aligned} \psi'(t) &= a_1 & \text{if } t \geq 0 \\ &= a_2 & \text{if } t < 0, \end{aligned}$$

where a_1 and a_2 are constants. Therefore, $\psi = \tilde{\psi}$ a.e. w.r.t. μ_1 where

$$\begin{aligned} \tilde{\psi}(x) &= a_1 x + b & \text{if } x \geq 0 \\ &= a_2 x + b & \text{if } x < 0, \end{aligned}$$

and b is a constant. But the equality $\text{Var}[\psi(X)] = E[\psi'(X)]^2 < \infty$ also implies that

$$\begin{aligned} \text{Var}[\tilde{\psi}(X)] &= \text{Var}[\psi(X)] = E \left[\int_0^x \psi'(t) dt \right]^2 \\ &= E[\tilde{\psi}(X) - \tilde{\psi}(0)]^2. \end{aligned}$$

Therefore, $E\tilde{\psi}(X) = \tilde{\psi}(0) = b$. Consequently, $a_1 = a_2$, and we have $\psi(x) = ax + b$ a.e. w.r.t. μ_1 for some constants a and b . This completes the proof of the lemma.

THEOREM 2.1. *Let X_1, \dots, X_k be independent random variables each having the standard normal distribution. Let g, g_1, \dots, g_k be real-valued Borel measurable functions defined on \mathbb{R}^k such that*

$$\begin{aligned} g(x_1, \dots, x_k) &= \int_0^{x_i} g_i(x_1, \dots, x_{i-1}, t, x_{i+1}, \dots, x_k) dt \\ &\quad + g(x_1, \dots, x_{i-1}, 0, x_{i+1}, \dots, x_k) \end{aligned} \quad (2.1)$$

a.e. w.r.t. μ_k for $i = 1, \dots, k$. Then

$$\text{Var}[g(X_1, \dots, X_k)] \leq \sum_{i=1}^k E[g_i(X_1, \dots, X_k)]^2, \quad (2.2)$$

where equality holds with the right-hand side finite if and only if $g(x_1, \dots, x_k) = a_1 x_1 + \dots + a_k x_k + b$ a.e. w.r.t. μ_k for some constants a_1, \dots, a_k and b .

Proof. If $k = 1$, the theorem follows trivially from Lemma 2.1. So let $k \geq 2$. There is nothing to prove if $\sum_{i=1}^k E[g_i(X_1, \dots, X_k)]^2 = \infty$. So we assume it to be finite. The first thing to do is to show that

$$E[g(X_1, \dots, X_k)]^2 < \infty. \quad (2.3)$$

We do this by induction. By Fubini's theorem,

$$E[g_i(x_1, \dots, x_{i-1}, X_i, \dots, X_k)]^2 < \infty \quad (2.4)$$

for almost all (x_1, \dots, x_{i-1}) w.r.t. μ_{i-1} and for $i = 1, \dots, k$. Here and throughout the rest of the proof, the statement "for almost all (x_1, \dots, x_{i-1}) w.r.t. μ_{i-1} " is to be omitted for $i = 1$. By Lemma 2.1 and the independence of X_1, \dots, X_k we have

$$\begin{aligned} E \left[\int_0^{X_i} g_i(x_1, \dots, x_{i-1}, t, X_{i+1}, \dots, X_k) dt \right]^2 \\ \leq E[g_i(x_1, \dots, x_{i-1}, X_i, \dots, X_k)]^2 \end{aligned} \quad (2.5)$$

for almost all (x_1, \dots, x_{i-1}) w.r.t. μ_{i-1} and for $i = 1, \dots, k$. We start with $i = k$. From condition (2.1), we have, for almost all (x_1, \dots, x_{k-1}) w.r.t. μ_{k-1} ,

$$\begin{aligned} g(x_1, \dots, x_{k-1}, X_k) &= g(x_1, \dots, x_{k-1}, 0) \\ &+ \int_0^{X_k} g_k(x_1, \dots, x_{k-1}, t) dt \quad \text{a.s.} \end{aligned}$$

Combining this with (2.5) we obtain

$$E[g(x_1, \dots, x_{k-1}, X_k)]^2 < \infty$$

for almost all (x_1, \dots, x_{k-1}) w.r.t. μ_{k-1} . Now assume that

$$E[g(x_1, \dots, x_i, X_{i+1}, \dots, X_k)]^2 < \infty \quad (2.6)$$

for almost all (x_1, \dots, x_i) w.r.t. μ_i . Since the univariate normal density function is bounded away from zero on any bounded interval, it follows from (2.4) that

$$E \left| \int_0^{x_i} [g_i(x_1, \dots, x_{i-1}, t, X_{i+1}, \dots, X_k)]^2 dt \right| < \infty \quad (2.7)$$

for almost all (x_1, \dots, x_{i-1}) w.r.t. μ_{i-1} and all x_i . This together with (2.1) and (2.6) imply that

$$E[g(x_1, \dots, x_{i-1}, 0, X_{i+1}, \dots, X_k)]^2 < \infty \quad (2.8)$$

for almost all (x_1, \dots, x_{i-1}) w.r.t. μ_{i-1} . Combining (2.1), (2.5) and (2.8), we obtain

$$E[g(x_1, \dots, x_{i-1}, X_i, \dots, X_k)]^2 < \infty$$

for almost all (x_1, \dots, x_{i-1}) w.r.t. μ_{i-1} . This proves (2.3).

Now let \mathcal{F}_0 be the trivial σ -algebra and \mathcal{F}_i be the σ -algebra generated by X_1, \dots, X_i , $i = 1, \dots, k$. Write E_i for the conditional expectation operator given \mathcal{F}_i , $i = 0, 1, \dots, k$. Define $Y_i = E_i g(X_1, \dots, X_k) - E_{i-1} g(X_1, \dots, X_k)$ for $i = 1, \dots, k$. Then $\{Y_i : i = 1, \dots, k\}$ is a sequence of martingale differences relative to $\{\mathcal{F}_i : i = 1, \dots, k\}$ and $g(X_1, \dots, X_k) - Eg(X_1, \dots, X_k) = \sum_{i=1}^k Y_i$. By the martingale property,

$$\text{Var}[g(X_1, \dots, X_k)] = \sum_{i=1}^k EY_i^2. \quad (2.9)$$

But by the independence of X_1, \dots, X_k , Eq. (2.1) and Lemma 2.1, we have for $i = 1, \dots, k$,

$$\begin{aligned} EY_i^2 &= E \left[E_i \int_0^{X_i} g_i(X_1, \dots, X_{i-1}, t, X_{i+1}, \dots, X_k) dt \right. \\ &\quad \left. - E_{i-1} \int_0^{X_i} g_i(X_1, \dots, X_{i-1}, t, X_{i+1}, \dots, X_k) dt \right]^2 \\ &\leq E \left[E_i \int_0^{X_i} g_i(X_1, \dots, X_{i-1}, t, X_{i+1}, \dots, X_k) dt \right]^2 \\ &= EE_{i-1} \left[\int_0^{X_i} E_i g_i(X_1, \dots, X_{i-1}, t, X_{i+1}, \dots, X_k) dt \right]^2 \\ &\leq EE_{i-1} [E_i g_i(X_1, \dots, X_k)]^2 \\ &\leq E[g_i(X_1, \dots, X_k)]^2. \end{aligned} \quad (2.10)$$

This proves (2.2).

If $g(x_1, \dots, x_k) = a_1 x_1 + \dots + a_k x_k + b$ a.e. w.r.t. μ_k , then inequality (2.2) trivially reduces to an equality with both sides finite. Conversely, if equality in (2.2) holds with the right-hand side finite, then by (2.9) and (2.10), $EY_i^2 = E[g_i(X_1, \dots, X_k)]^2 < \infty$ for $i = 1, \dots, k$, which, in turn, implies that

$$\begin{aligned} \text{Var}_{i-1} \left[\int_0^{X_i} E_i g_i(X_1, \dots, X_{i-1}, t, X_{i+1}, \dots, X_k) dt \right]^2 \\ = E_{i-1} [E_i g_i(X_1, \dots, X_k)]^2 \quad \text{a.s.}, \end{aligned}$$

where Var_{i-1} denotes conditional variance given \mathcal{F}_{i-1} . By Lemma 2.1,

$$\int_0^{X_i} E_i g_i(X_1, \dots, X_{i-1}, t, X_{i+1}, \dots, X_k) dt = u_i X_i + v_i \quad \text{a.s.},$$

where u_i and v_i are \mathcal{F}_{i-1} measurable, $i = 1, \dots, k$. It follows that for $i = 1, \dots, k$,

$$Y_i = u_i X_i \quad \text{a.s.}$$

We now prove by induction that all the u_i must be constants a.s. We have proved that u_1 is a constant a.s. Suppose u_1, \dots, u_j are constants a.s., say $u_1 = a_1, \dots, u_j = a_j$ a.s., $1 \leq j < k$. Then

$$\begin{aligned} g(X_1, \dots, X_k) - E g(X_1, \dots, X_k) \\ = a_1 X_1 + \dots + a_j X_j + u_{j+1} X_{j+1} + \dots + u_k X_k \quad \text{a.s.} \quad (2.11) \end{aligned}$$

Now rearrange X_1, \dots, X_k in the order $X_{j+1}, X_1, \dots, X_j, X_{j+2}, \dots, X_k$. Define $\tilde{\mathcal{F}}_i, \tilde{E}_i, \tilde{Y}_i$ with respect to $X_{j+1}, X_1, \dots, X_j, X_{j+2}, \dots, X_k$ in the same way as \mathcal{F}_i, E_i, Y_i are with respect to X_1, \dots, X_k , $i = 1, \dots, k$. For example, $\tilde{\mathcal{F}}_1, \tilde{\mathcal{F}}_2$, etc., are respectively σ -algebras generated by $\{X_{j+1}\}$, $\{X_{j+1}, X_1\}$, etc. Then

$$\begin{aligned} \tilde{Y}_1 &= \tilde{E}_1(a_1 X_1 + \dots + a_j X_j + u_{j+1} X_{j+1} + \dots + u_k X_k) \\ &= E(a_1 X_1 + \dots + a_j X_j + u_{j+1} X_{j+1} + \dots + u_k X_k \mid X_{j+1}) \\ &= (E u_{j+1}) X_{j+1} \quad \text{a.s.} \end{aligned}$$

and

$$E(\tilde{Y}_1)^2 = (E u_{j+1})^2.$$

But by arguments similar to those leading to (2.9) and (2.10), the equality in (2.2) implies that

$$\begin{aligned} E(\tilde{Y}_1)^2 &= E[g_{j+1}(X_1, \dots, X_k)]^2 \\ &= E Y_{j+1}^2 = E u_{j+1}^2. \end{aligned}$$

It follows that

$$Eu_{j+1}^2 = (Eu_{j+1})^2$$

which is $\text{Var}(u_{j+1}) = 0$ and hence implies that u_{j+1} is a constant a.s. This proves that all the u_i are constants a.s. Consequently,

$$g(x_1, \dots, x_k) = a_1 x_1 + \dots + a_k x_k + b \quad \text{a.e. w.r.t. } \mu_k,$$

where a_1, \dots, a_k and $b (=Eg(X_1, \dots, X_k))$ are constants. This completes the proof of the theorem.

3. COROLLARIES

In this section we consider inequality (2.2) for two specific classes of functions which satisfy (2.1).

COROLLARY 3.1. *Let X_1, \dots, X_k be independent random variables each having the standard normal distribution. Let g be a partially differentiable real-valued function defined on \mathbb{R}^k . Then*

$$\text{Var}[g(X_1, \dots, X_k)] \leq \sum_{i=1}^k E[D_i g(X_1, \dots, X_k)]^2,$$

where equality holds with the right-hand side finite if and only if

$$g(x_1, \dots, x_k) = a_1 x_1 + \dots + a_k x_k + b$$

for all x_1, \dots, x_k , and some constants a_1, \dots, a_k and b .

We note that partially differentiable functions are separately continuous (that is, continuous in each variable) and that separately continuous functions from \mathbb{R}^k to \mathbb{R} are Borel measurable (see, for example, Kuratowski [3, p. 285]). The proof of Corollary 3.1 follows trivially from these facts and the following lemma.

LEMMA 3.1. *Let $h: \mathbb{R}^k \rightarrow \mathbb{R}$ be separately continuous. If $h = 0$ a.e. w.r.t. μ_k , then $h = 0$ everywhere.*

Proof. Suppose there is a point at which $h \neq 0$. Without loss of generality, we may assume that $h(0, \dots, 0) = 2$. By the separate continuity of h , there exist an open interval J_1 containing 0 such that

$$h(x_1, 0, \dots, 0) > 1 \quad \text{for } x_1 \in J_1.$$

For each $x_1 \in J_1$, there exists an open interval $J_2(x_1)$ containing 0 such that

$$h(x_1, x_2, 0, \dots, 0) > \frac{1}{2} \quad \text{for } x_2 \in J_2(x_1).$$

Arguing inductively, we have

$$h(x_1, \dots, x_k) > \frac{1}{k}.$$

for $x_i \in J_i(x_1, \dots, x_{i-1})$ for $i = 1, \dots, k$ where each $J_i(x_1, \dots, x_{i-1})$ is an open interval containing 0 and $J_i(x_1, \dots, x_{i-1})$ is defined to be J_1 for $i = 1$. Let $A = \{(x_1, \dots, x_k): x_i \in J_i(x_1, \dots, x_{i-1}), i = 1, \dots, k\}$. Clearly $A \subset \{(x_1, \dots, x_k): h(x_1, \dots, x_k) > 1/k\}$. But $h = 0$ a.e. w.r.t. μ_k . This implies that $\mu_k(h > 1/k) = 0$ which, in turn, implies that A is Lebesgue measurable and $\mu_k(A) = 0$. Now define

$$A(n_2, \dots, n_k) = \left\{ (x_1, \dots, x_k) \in A: \mu_1(J_j(x_1, \dots, x_{j-1})) > \frac{1}{n_j}, j = 2, \dots, k \right\}.$$

Then $A = \bigcup_{n_2=1}^{\infty} \dots \bigcup_{n_k=1}^{\infty} A(n_2, \dots, n_k)$ so that for each (n_2, \dots, n_k) , $A(n_2, \dots, n_k)$ is Lebesgue measurable and $\mu_k(A(n_2, \dots, n_k)) = 0$. Define

$$B(n_2, \dots, n_k) = \{x_1: (x_1, \dots, x_k) \in A(n_2, \dots, n_k)\}.$$

Clearly $J_1 = \bigcup_{n_2=1}^{\infty} \dots \bigcup_{n_k=1}^{\infty} B(n_2, \dots, n_k)$ and $B(n_2, \dots, n_k)$ is nondecreasing in n_2, \dots, n_k . Therefore, for sufficiently large n_2, \dots, n_k , $\mu_1(B(n_2, \dots, n_k)) \geq \frac{1}{2}\mu_1(J_1) > 0$. But for sufficiently large n_2, \dots, n_k , we have by Fubini's theorem

$$\begin{aligned} 0 = \mu_k(A(n_2, \dots, n_k)) &= \int \dots \int_{(x_1, \dots, x_k) \in A(n_2, \dots, n_k)} d\mu_1(x_k) \dots d\mu_1(x_1) \\ &\geq \frac{1}{n_2 \dots n_k} \int_{x_1 \in B(n_2, \dots, n_k)} d\mu_1(x_1) \\ &\geq \frac{\mu_1(J_1)}{2n_2 \dots n_k} > 0 \end{aligned}$$

which is a contradiction. This proves the lemma.

If g has a differential at every point in \mathbb{R}^k , an inequality for the multivariate normal distribution with zero mean vector and any covariance matrix Σ can be deduced from Corollary 3.1. For the next corollary, let m ($1 \leq m \leq k$) be the rank of Σ and $\mathcal{L}(\Sigma)$ denote the real linear space spanned by the column vectors of Σ . Let v_1, \dots, v_m be column vectors which form a basis of $\mathcal{L}(\Sigma)$ and g_{Σ} be the restriction of g to $\mathcal{L}(\Sigma)$. The gradient of g will

be denoted by ∇g and be represented by a column vector. The transpose of a matrix A will be denoted by A^t .

COROLLARY 3.2. *Let $\xi = (X_1, \dots, X_k)^t$ be a random vector having the multivariate normal distribution with zero mean vector and covariance matrix Σ . Let g be a real-valued function defined on \mathbb{R}^k such that it has a differential at every point. Then*

$$\text{Var}[g(\xi)] \leq E \nabla^t g(\xi) \Sigma \nabla g(\xi), \quad (3.1)$$

where equality holds with the right-hand side finite if and only if $g_\Sigma(x_1 v_1 + \dots + x_m v_m) = a_1 x_1 + \dots + a_m x_m + b$ for all real numbers x_1, \dots, x_m and some constants a_1, \dots, a_m and b .

Proof. We note that the form of g_Σ corresponding to equality in (3.1) is independent of the choice of the basis $\{v_1, \dots, v_m\}$ of $\mathcal{L}(\Sigma)$. Thus, we shall choose v_1, \dots, v_m to be such that $\xi = Y_1 v_1 + \dots + Y_m v_m$ with Y_1, \dots, Y_m being independent random variables each having the standard normal distribution. Let v_{m+1}, \dots, v_k be linearly independent column vectors such that $\{v_1, \dots, v_k\}$ is a basis of \mathbb{R}^k . Let $C = (v_1, \dots, v_m)$ and $A = (v_1, \dots, v_k) = (c_{ij})$. Then $\Sigma = CC^t$. Consider the mapping $f: \mathbb{R}^k \rightarrow \mathbb{R}^k$ defined by $f(y) = Ay$, $y \in \mathbb{R}^k$. Define $h: \mathbb{R}^k \rightarrow \mathbb{R}$ by $h = g \circ f$. Since g has a differential, so has h and

$$\nabla h(y) = \sum_{i=1}^k D_i g(f(y))(c_{i1}, \dots, c_{ik})^t.$$

Therefore,

$$\begin{aligned} \nabla^t g(\xi) \Sigma \nabla g(\xi) &= \nabla^t g(\xi) CC^t \nabla g(\xi) \\ &= \sum_{j=1}^m [v_j^t \nabla g(\xi)]^2 = \sum_{j=1}^m \left[\sum_{i=1}^k c_{ij} D_i g(\xi) \right]^2 \\ &= \sum_{j=1}^m [D_j h(Y_1, \dots, Y_m, 0, \dots, 0)]^2. \end{aligned} \quad (3.2)$$

There is nothing to prove if $E \nabla^t g(\xi) \Sigma \nabla g(\xi) = \infty$. However, if $E \nabla^t g(\xi) \Sigma \nabla g(\xi) < \infty$, then, by (3.2), $\sum_{j=1}^m E [D_j h(Y_1, \dots, Y_m, 0, \dots, 0)]^2 < \infty$. Applying Corollary 3.1, we obtain

$$\begin{aligned} \text{Var}[g(\xi)] &= \text{Var}[h(Y_1, \dots, Y_m, 0, \dots, 0)] \\ &\leq \sum_{j=1}^m E [D_j h(Y_1, \dots, Y_m, 0, \dots, 0)]^2 \\ &= E \nabla^t g(\xi) \Sigma \nabla g(\xi). \end{aligned}$$

It is easy to check that if $g_{\Sigma}(x_1 v_1 + \cdots + x_m v_m) = a_1 x_1 + \cdots + a_m x_m + b$ for some constants a_1, \dots, a_m and b , then (3.1) reduces to an equality with the right-hand side finite. Conversely, if the latter holds, then

$$\text{Var}[h(Y_1, \dots, Y_m, 0, \dots, 0)] = \sum_{j=1}^m E[D_j h(Y_1, \dots, Y_m, 0, \dots, 0)]^2$$

and by Corollary 3.1 again

$$h(x_1, \dots, x_m, 0, \dots, 0) = a_1 x_1 + \cdots + a_m x_m + b$$

for some constants a_1, \dots, a_m and b . But

$$h(x_1, \dots, x_m, 0, \dots, 0) = g_{\Sigma}(x_1 v_1 + \cdots + x_m v_m).$$

This completes the proof of the corollary.

ACKNOWLEDGMENTS

The author is thankful to Dr. Huei Chuen Huang for a discussion which led to the proof of Lemma 3.1. He is also thankful to the referee for his suggestions which helped to simplify the proofs of Lemma 2.1 and Theorem 2.1.

REFERENCES

- [1] CHERNOFF, H. (1980). The identification of an element of a large population in the presence of noise. *Ann. Statist.* **8** 1179–1197.
- [2] CHERNOFF, H. (1981). A note on an inequality involving the normal distribution. *Ann. Probab.* **9** 533–535.
- [3] KURATOWSKI, C. (1958). *Topologie*. Panstwowe Wydawnictwo Naukowe, Warszawa.